
Children's Hospital and Health System Chorus Community Health Plans Policy and Procedure



Medical Utilization Management Policy

SUBJECT: CUSTOM FOOT ORTHOTICS

INCLUDED PRODUCT(S):

| Medicaid | Individual and Family |
|---------------------|-----------------------|
| BadgerCare Plus | Commercial |
| 🛛 Care4Kids Program | 🛛 Marketplace |

PURPOSE OR DESCRIPTION:

The purpose of this policy is to define criteria for the medically necessary use of custom foot orthotics (CFO).

POLICY:

CCHP considers a custom foot orthoic (CFO) to be medically necessary when a member has pain in the arch, heel or other area of the foot and diagnosis of:

- 1. Juvenile idiopathic arthritis
- 2. Pes cavus (high arch)
- 3. Other, including pes planus, pes planovalgus and plantar fasciitis:
 - a. Symptoms present for at least 3 months
 - b. Conservative measures have been tried first, as indicated by one of the following :
 - i. Physical therapy
 - ii. Prefabricated orthotics

Effective: 5/16 Last reviewed: 9/23 Q: \CCHP Leadership\Utilization Management Medical Policies\APPROVED MEDICAL UM POLICIES\Custom Foot Orthotics Medical UM Policy Developed by: CCHP Medical Directors

- iii. Anti-inflammatory medication
- iv. Activity adjustment
- c. Member has been evaluated by the prescribing provider AFTER the conservative measure(s) has been tried, and the ongoing need for the CFO is documented.

For CFO replacement, ALL of the following must be present:

- 1. At least 12 months have passed since the last CFO was dispensed
- 2. Documentation shows one of the following:
 - a. The initial CFO is worn out; or
 - b. The member is a growing child, and the initial CFO is too small
- 3. Documentation shows one of the following:
 - a. The initial CFO was effective and member was using the CFO; or
 - b. Member has been re-evaluated by the prescribing provider and documentation supports the ongoing need for the CFO

REFERENCES

- 1. Anain JM, Bojrab AR, Rhinehart FC. Conservative treatments for rheumatoid arthritis in the foot and ankle. Clinics in Podiatric Medicine and Surgery 2010;27(2):193-207. DOI: 10.1016/j.cpm.2009.12.003.
- 2. Anderson J, Stanek J. Effect of foot orthoses as treatment for plantar fasciitis or heel pain. Journal of Sport Rehabilitation 2013;22(2):130-6.
- 3. Baker K, et al. A randomized crossover trial of a wedged insole for treatment of knee osteoarthritis. Arthritis and Rheumatism 2007;56(4):1198-203. DOI: 10.1002/art.22516.
- 4. Baldassin V, Gomes CR, Beraldo PS. Effectiveness of prefabricated and customized foot orthoses made from low-cost foam for noncomplicated plantar fasciitis: a randomized controlled trial. Archives of Physical Medicine and Rehabilitation 2009;90(4):701-6. DOI: 10.1016/j.apmr.2008.11.002.
- Burns J, Landorf KB, Ryan MM, Crosbie J, Ouvrier RA. Interventions for the prevention and treatment of pes cavus. Cochrane Database of Systematic Reviews 2007, (verified by Cochrane 2010 Dec), Issue 4. Art. No.: CD006154. DOI: 10.1002/14651858.CD006154.pub2.
- Cameron-Fiddes V, Santos D. The use of 'off-the-shelf' foot orthoses in the reduction of foot symptoms in patients with early rheumatoid arthritis. Foot (Edinburgh, Scotland) 2013;23(4):123-9. DOI: 10.1016/j.foot.2013.09.001.
- Coda A, Fowlie PW, Davidson JE, Walsh J, Carline T, Santos D. Foot orthoses in children with juvenile idiopathic arthritis: a randomised controlled trial. Archives of Disease in Childhood 2014;99(7):649-51. DOI: 10.1136/archdischild-2013-305166.
- 8. Covey CJ, Mulder MD. Plantar fasciitis: How best to treat? Journal of Family Practice 2013;62(9):466-71.
- 9. Cutts S, Obi N, Pasapula C, Chan W. Plantar fasciitis. Annals of the Royal College of Surgeons of England 2012;94(8):539-42. DOI: 10.1308/003588412X13171221592456.
- DiGiovanni BF, Moore AM, Zlotnicki JP, Pinney SJ. Preferred management of recalcitrant plantar fasciitis among orthopaedic foot and ankle surgeons. Foot and Ankle International 2012;33(6):507-12. DOI: 10.3113/FAI.2012.0507.
- 11. Drake M, Bittenbender C, Boyles RE. The short-term effects of treating plantar fasciitis with a temporary custom foot orthosis and stretching. Journal of Orthopaedic and Sports Physical Therapy 2011;41(4):221-31. DOI: 10.2519/jospt.2011.3348.
- 12. Evans AM, Rome K. A Cochrane review of the evidence for non-surgical interventions for flexible pediatric flat feet. European Journal of Physical and Rehabilitation Medicine 2011;47(1):69-89.
- 13. Gelis A, Coudeyre E, Hudry C, Pelissier J, Revel M, Rannou F. Is there an evidence-based efficacy for the use of foot orthotics in knee and hip osteoarthritis? Elaboration of French clinical practice guidelines. Joint, Bone, Spine 2008;75(6):714-20. DOI: 10.1016/j.jbspin.2008.02.013.

- 14. Hawke F, Burns J. Brief report: custom foot orthoses for foot pain: what does the evidence say? Foot and Ankle International 2012;33(12):1161-3. DOI: DOI: 10.3113/FAI.2012.1161.
- Hawke F, Burns J, Radford JA, du Toit V. Custom-made foot orthoses for the treatment of foot pain. Cochrane Database of Systematic Reviews 2008, (verified by Cochrane 2008 Mar), Issue 3. Art. No.: CD006801. DOI: 10.1002/14651858.CD006801.pub2.
- Hendry GJ, Turner DE, Lorgelly PK, Woodburn J. Room for improvement: patient, parent, and practitioners' perceptions of foot problems and foot care in juvenile idiopathic arthritis. Archives of Physical Medicine and Rehabilitation 2012;93(11):2062-7. DOI: 10.1016/j.apmr.2012.07.007.
- 17. Hennessy K, Woodburn J, Steultjens MP. Custom foot orthoses for rheumatoid arthritis: A systematic review. Arthritis Care & Research 2012;64(3):311-20. DOI: 10.1002/acr.21559.
- Lee SY, McKeon P, Hertel J. Does the use of orthoses improve self-reported pain and function measures in patients with plantar fasciitis? A meta-analysis. Physical Therapy in Sport 2009;10(1):12-8. DOI: 10.1016/j.ptsp.2008.09.002.
- MacKenzie AJ, Rome K, Evans AM. The efficacy of nonsurgical interventions for pediatric flexible flat foot: a critical review. Journal of Pediatric Orthopedics 2012;32(8):830-4. DOI: 10.1097/BPO.0b013e3182648c95.
- Najafi B, Barnica E, Wrobel JS, Burns J. Dynamic plantar loading index: understanding the benefit of custom foot orthoses for painful pes cavus. Journal of Biomechanics 2012;45(9):1705-11. DOI: 10.1016/j.jbiomech.2012.03.006.
- 21. Oldfield V, Felson DT. Exercise therapy and orthotic devices in rheumatoid arthritis: evidence-based review. Current Opinion in Rheumatology 2008;20(3):353-9. DOI: 10.1097/BOR.0b013e3282fd17df.
- 22. Orchard J. Plantar fasciitis. British Medical Journal 2012;345:e6603.
- Parikh SS, Baxi N, Padavan SA. Musculoskeletal medicine. In: Sackheim KA, editor. Rehab Clinical Pocket Guide. New York, NY: Springer; 2013:357-426.
- Parkes MJ, et al. Lateral wedge insoles as a conservative treatment for pain in patients with medial knee osteoarthritis: a meta-analysis. Journal of the American Medical Association 2013;310(7):722-30. DOI: 10.1001/jama.2013.243229.
- Rao S, Riskowski JL, Hannan MT. Musculoskeletal conditions of the foot and ankle: assessments and treatment options. Best Practice and Research. Clinical Rheumatology 2012;26(3):345-68. DOI: 10.1016/j.berh.2012.05.009.
- Reilly KA, Barker KL, Shamley D. A systematic review of lateral wedge orthotics--how useful are they in the management of medial compartment osteoarthritis? Knee 2006;13(3):177-83. DOI: 10.1016/j.knee.2006.02.003.
- Rome K, Ashford RL, Evans A. Non-surgical interventions for paediatric pes planus. Cochrane Database of Systematic Reviews 2010, (verified by Cochrane 2012 Nov), Issue 7. Art. No.: CD006311. DOI: 10.1002/14651858.CD006311.pub2.
- Rosenbloom KB. Pathology-designed custom molded foot orthoses. Clinics in Podiatric Medicine and Surgery 2011;28(1):171-87. DOI: 10.1016/j.cpm.2010.11.001.
- 29. Van Lunen B, Cortes N, Andrus T, Walker M, Pasquale M, Onate J. Immediate effects of a heel-pain orthosis and an augmented low-dye taping on plantar pressures and pain in subjects with plantar fasciitis. Clinical Journal of Sport Medicine 2011;21(6):474-9. DOI: 10.1097/JSM.0b013e3182340199.
- 30. Vliet Vlieland TP, van den Ende CH. Nonpharmacological treatment of rheumatoid arthritis. Current Opinion in Rheumatology 2011;23(3):259-64. DOI: 10.1097/BOR.0b013e32834540fb.
- 31. Walther M, et al. Effect of different orthotic concepts as first line treatment of plantar fasciitis. Foot and Ankle Surgery 2013;19(2):103-7. DOI: 10.1016/j.fas.2012.12.008.