

Psychotropic Medication Utilization Parameters for Children and Youth in Foster Care (5th Version)



Developed by:

*Texas Department of Family and Protective Services
and*

The University of Texas at Austin College of Pharmacy

with review and input provided by:

- ❖ *Federation of Texas Psychiatry*
- ❖ *Texas Pediatric Society*
- ❖ *Texas Academy of Family Physicians*
- ❖ *Texas Medical Association*

and

*Rutgers University-Center for Education and Research
on Mental Health Therapeutics*

March 2016

(Tables Updated July 2016)

Table of Contents

❖ <i>Introduction and General Principles</i>	3-6
❖ <i>Criteria Indicating Need for Further Review of a Child's Clinical Status</i>	7
❖ <i>Psychotropic Medication Tables</i>	8-17
❖ <i>Glossary</i>	18
❖ <i>References</i>	19-22
❖ <i>Web Link References</i>	23
❖ <i>Members of the Ad Hoc Working Group</i>	24
❖ <i>Acknowledgements and Web Address for the Parameters Document</i>	25

Psychotropic Medication Utilization Parameters for Children and Youth in Foster Care

Introduction and General Principles

The use of psychotropic medications by children and youth is an issue confronting parents, other caregivers, and health care professionals across the United States. Children and youth in foster care, in particular, have multiple needs, including those related to emotional or psychological stress. They typically have experienced abusive, neglectful, serial or chaotic caretaking environments. Birth family history is often not available. These children often present with a fluidity of different symptoms over time reflective of past traumatic events that may mimic many psychiatric disorders and result in difficulties with attachment, mood regulation, behavioral control, and other areas of functioning.

Because of the complex issues involved in the lives of foster children, it is important that a comprehensive evaluation be performed before beginning treatment for a mental or behavioral disorder. Except in the case of an emergency, a child should receive a thorough health history, psychosocial assessment, mental status exam, and physical exam before prescribing a psychotropic medication. The physical assessment should be performed by a physician or another healthcare professional qualified to perform such an assessment. It is recognized that in some emergency situations, it may be in the best interest of the child to prescribe psychotropic medications before a physical exam can actually be performed. In these situations, a thorough health history should be performed to assess for significant medical disorders and past response to medications, and a physical evaluation should be performed as soon as possible. A thorough psychosocial assessment should be performed by an appropriately qualified mental health clinician

(masters or doctoral level), a psychiatrist/child psychiatrist, or a primary care physician with experience in providing mental health care to children and youth. The child's symptoms and functioning should be assessed across multiple domains, and the assessment should be developmentally age appropriate. It is very important that information about the child's history, including history of trauma and current functioning be made available to the treating physician in a timely manner, either through an adult who is well-informed about the child or through a comprehensive medical record. It is critical to meet the individual needs of patients and their families in a culturally competent manner. This indicates a need to address communication issues as well as differences in perspective on issues such as behavior and mental functioning. Interpretation of clinical symptoms and decisions concerning treatment should, whenever possible, be informed by the child's developmental history of trauma, neglect or abuse and the timing of these stressors. In general, optimal outcomes are achieved with well-coordinated team based care with members of different professions (e.g., child psychiatrist, child psychologist, social worker, primary care physician, etc.) each contributing their particular expertise to the treatment plan and follow-up. Additionally, at present there are no biomarkers to assist with the diagnosis of mental disorders, and imaging (e.g., MRI) and other tests (e.g., EEG) are not generally helpful in making a clinical diagnosis of a mental disorder.

The role of non-pharmacological interventions should be considered before beginning a psychotropic medication, except in urgent situations such as suicidal ideation, psychosis, self-injurious

behavior, physical aggression that is acutely dangerous to others, or severe impulsivity endangering the child or others; when there is marked disturbance of psychophysiological functioning (such as profound sleep disturbance), or when the child shows marked anxiety, isolation, or withdrawal. Given the history of trauma, unusual stress and change in environmental circumstances associated with being a child in foster care, psychotherapy should generally begin before or concurrent with prescription of a psychotropic medication. Referral for trauma-informed, evidence-based psychotherapy should be considered when available and appropriate. Equally important, the role of the health care provider and the health care environment's potential to exacerbate a child's symptoms, given their respective trauma history, should be considered and minimized. Patient and caregiver education should be provided about the condition to be treated, treatment options (non-pharmacological and pharmacological), treatment expectations, and potential side effects that may occur during the prescription of psychotropic medications.

It is recognized that many psychotropic medications do not have Food and Drug Administration (FDA) approved labeling for use in children. The FDA has a statutory mandate to determine whether pharmaceutical company sponsored research indicates that a medication is safe and effective for those indications that are listed in the approved product labeling. The FDA assures that information in the approved product labeling is accurate, and limits the manufacturer's marketing to the information contained in the approved labeling. The FDA does not regulate physician and other health provider practice. In fact,

the FDA has stated that it does “not limit the manner in which a practitioner may prescribe an approved drug.” Studies and expert clinical experience often support the use of a medication for an “off-label” use. Physicians should utilize the available evidence, expert opinion, their own clinical experience, and exercise their clinical judgment in prescribing what is best for each individual patient. To that end, clear documentation of the physician’s rationale in the medical record facilitates continuity of care and minimizes misinterpretation.

Role of Primary Care Providers

Primary care providers play a valuable role in the care of youth with mental disorders. Not only are they the clinicians most likely to initially interact with children who are in distress due to an emotional or psychiatric disorder, but also an inadequate number of child psychiatrists are available to meet all children’s mental health needs. Primary care clinicians are in an excellent position to perform screenings of children for potential mental disorders, and they should be able to diagnose and treat relatively straightforward situations such as uncomplicated ADHD, anxiety, or depression. Primary care providers should provide advice to youth in foster care and their care givers about handling feelings and behaviors, recognizing the need for help, making decisions regarding healthy life styles, and the available treatments for childhood mental disorders. As always, consideration should be given regarding the need for referral for counseling, psychotherapy, or behavioral therapy. Primary care providers vary in their training, clinical experience, and confidence to address mental disorders in children. Short courses and intensive skills oriented seminars may be beneficial in assisting primary care clinicians in caring for children with mental disorders. Active liaisons with child psychiatrists who are available for phone consultation or referral can be beneficial in assisting primary care clinicians to meet the mental health needs of children. “The management of common presentations of ADHD, depression and anxiety, psychotherapy referral, psychopharmacology and appro-

priate child psychiatry referral are within the scope of general pediatric practice” (Southammakosane 2015). In addition, the American Academy of Pediatrics has recently provided a policy statement (“Health Care Issues for Children and Adolescents in Foster Care and Kinship Care”) which can be found at:

<http://pediatrics.aappublications.org/content/136/4/e1131>

General principles regarding the use of psychotropic medications in children include:

- A DSM-5 psychiatric diagnosis should be made before the prescribing of psychotropic medications.
- Clearly defined target symptoms and treatment goals for the use of psychotropic medications should be identified and documented in the medical record at the time of or before beginning treatment with a psychotropic medication. These target symptoms and treatment goals should be assessed at each clinic visit with the child and caregiver in a culturally and linguistically appropriate manner. Whenever possible, standardized clinical rating scales (clinician, patient, primary caregiver, teachers, and other care providers) or other measures should be used to quantify the response of the child’s target symptoms to treatment and the progress made toward treatment goals.
- In making a decision regarding whether to prescribe a psychotropic medication in a specific child, the clinician should carefully consider potential side effects, including those that are uncommon but potentially severe, and evaluate the overall benefit to risk ratio of pharmacotherapy.
- Except in the case of an emergency, informed consent should be obtained from the appropriate party(s) before beginning psychotropic medication. Informed consent to treatment with psychotropic medication entails diagnosis, expected benefits and risks of

treatment, including common side effects, discussion of laboratory findings, and uncommon but potentially severe adverse events. Alternative treatments, the risks associated with no treatment, and the overall potential benefit to risk ratio of treatment should be discussed.

- Whenever possible, trauma-informed, evidence-based psychotherapy, should begin before or concurrent with the prescription of psychotropic medication.
- Before starting psychopharmacological treatment in preschool-aged children even more emphasis should be placed on treatment with non-psychopharmacological interventions. Assessment of parent functioning and mental health needs, in addition to training parents in evidence-based behavior management can also reduce the need for the use of medication.
- Medication management should be collaborative. Youth, as well as caregivers, should be involved in decision-making about treatment, in accordance with their developmental level. Parents providing informed consent should be engaged, and where applicable, other caregivers, family, and child related agencies should be involved.
- During the prescription of psychotropic medication, the presence or absence of medication side effects should be documented in the child’s medical record at each visit.
- Appropriate monitoring of indices such as height, weight, blood pressure, or laboratory findings should be documented.
- Monotherapy regimens for a given disorder or specific target symptoms should usually be tried before polypharmacy regimens. While the goal is to use as few psychotropic medications as can be used to appropriately address the child’s clinical status, it is recognized that the presence of psychiatric comorbidities may affect the number of

psychotropic medications that are prescribed. When polypharmacy regimens are needed, addition of medications should occur in a systematic orderly process, accompanied by on-going monitoring, evaluation, and documentation. The goal remains to minimize polypharmacy while maximizing therapeutic outcomes.

- Medications should be initiated at the lower end of the recommended dose range and titrated carefully as needed.
- Only one medication should be changed at a time, unless a clinically appropriate reason to do otherwise is documented in the medical record. (Note: starting a new medication and beginning the dose taper of a current medication is considered one medication change).
- The use of “prn” or as needed prescriptions is discouraged. If they are used, the situation indicating need for the administration of a prn medication should be clearly indicated as well as the maximum dosage in a 24 hour period and in a week. The frequency of administration should be monitored to assure that these do not become regularly scheduled medications unless clinically indicated.
- The frequency of clinician follow-up should be appropriate for the severity of the child’s condition and adequate to monitor response to treatment, including: symptoms, behavior, function, and potential medication side effects. At a minimum, a child receiving psychotropic medication should be seen by the clinician at least once every ninety days.
- The potential for emergent suicidality should be carefully evaluated and monitored, particularly in depressed children and adolescents as well as those initiating antidepressants, those having a history of suicidal behavior or deliberate self-harm and those with a history of anxiety or substance abuse disorders.
- If the prescribing clinician is not a child psychiatrist, referral to or consultation with a child psychiatrist, or a general psychiatrist with significant experience in treating children, should occur if the child’s clinical status has not shown meaningful improvement within a timeframe that is appropriate for the child’s diagnosis and the medication regimen being used.
- Before adding additional psychotropic medications to a regimen, the child should be assessed for adequate medication adherence, appropriateness of medication daily dosage, accuracy of the diagnosis, the occurrence of comorbid disorders (including substance abuse and general medical disorders), and the influence of psychosocial stressors.
- If a medication has not resulted in improvement in a child’s target symptoms (or rating scale score), discontinue that medication rather than adding a second medication to it.
- If a medication is being used in a child for a primary target symptom of aggression associated with a DSM-5 non-psychotic diagnosis (e.g., conduct disorder, oppositional defiant disorder, intermittent explosive disorder), and the behavior disturbance has been in remission for six months, then serious consideration should be given to slow tapering and discontinuation of the medication. If the medication is continued in this situation, the necessity for continued treatment should be evaluated and documented in the medical record at a minimum of every six months.
- The clinician should clearly document care provided in the child’s medical record, including history, mental status assessment, physical findings (when relevant), impressions, rationale for medications prescribed, adequate laboratory monitoring specific to the drug(s) prescribed at intervals required specific to the prescribed drug and potential known risks, medication

response, presence or absence of side effects, treatment plan, and intended use of prescribed medications.

Use of Psychotropic Medication in Preschool Age Children

The use of psychotropic medication in young children of preschool ages is a practice that is limited by the lack of evidence available for use of these agents in this age group. The Preschool Psychopharmacology Working Group (PPWG) published guidelines (Gleason 2007) summarizing available evidence for use of psychotropic medications in this age group. The PPWG was established in response to the clinical needs of preschoolers being treated with psychopharmacological agents and the absence of systematic practice guidelines for this age group, with its central purpose to attempt to promote an evidence-based, informed, and clinically sound approach when considering medications in preschool-aged children.

The PPWG guidelines emphasize consideration of multiple different factors when deciding on whether to prescribe psychotropic medications to preschool-aged children. Such factors include the assessment and diagnostic methods utilized in evaluating the child for psychiatric symptoms/illness, the current state of knowledge regarding the impact of psychotropic medication use on childhood neurodevelopmental processes, the regulatory and ethical contexts of use of psychotropic medications in small children (including available safety information and FDA status), and the existing evidence base for use of psychotropic medication in preschool aged children.

The publication includes specific guidelines and algorithm schematics developed by the PPWG to help guide treatment decisions for a number of psychiatric disorders that may present in preschool-aged children, including Attention-Deficit Hyperactivity Disorder, Disruptive Behavioral Disorders, Major Depressive Disorder, Bipolar Disorder, Anxiety Disorders, Post-Traumatic Stress Disorder, Obsessive-Compulsive Disorder, Pervasive Developmental

Disorders, and Primary Sleep Disorders. The working group's key points and guidelines are similar to the general principles regarding the use of psychotropic medication in children already detailed in this paper. However, the working group's algorithms put more emphasis on treating preschool-aged children with non-pharmacological interventions (for up to 12 weeks) before starting psychopharmacological treatment, in an effort to be very cautious in introducing psychopharmacological interventions to rapidly developing preschoolers.

The working group also emphasizes the need to assess parent functioning and mental health needs, in addition to training parents in evidence-based behavior management, since parent behavior and functioning can have a large impact on behavior and symptoms in preschool-aged children.

Distinguishing between Levels of Warnings Associated with Medication Adverse Effects

Psychotropic medications have the potential for adverse effects, some that are treatment limiting. Some adverse effects are detected prior to marketing, and are included in the FDA approved product labeling provided by the manufacturers. When looking at product labeling, these adverse effects will be listed in the "Warnings and Precautions" section. As well, the "Adverse Reactions" section of the product labeling will outline those adverse effects reported during clinical trials, as well as those discovered during post-marketing evaluation. Many tertiary drug information resources also list common adverse effects and precautions for use with psychotropic medications.

At times, post-marketing evaluation may detect critical adverse effects associated with significant morbidity and mortality. The Food and Drug Administration (FDA) may require manufacturers to revise product labeling to indicate these critical adverse effects. If found to be particularly signifi-

cant, these effects are demarcated by a box outlining the information at the very beginning of the product labeling, and have, in turn, been named boxed warnings. Boxed warnings are the strongest warning required by the FDA. It is important for clinicians to be familiar with all medication adverse effects, including boxed warnings, in order to appropriately monitor patients and minimize the risk of their occurrence. The medication tables include boxed warnings as well as other potential adverse effects. The list of potential adverse effects in the tables should not be considered exhaustive, and the clinician should consult the FDA approved product labeling and other reliable sources for information regarding medication adverse effects.

The FDA has in recent years taken additional measures to try to help patients avoid serious adverse events. New guides called Medication Guides have been developed, and are specific to particular medication and medication classes. Medication Guides advise patients and caregivers regarding possible adverse effects associated with classes of medications, and include precautions that they or healthcare providers may take while taking/prescribing certain classes of medications. The FDA requires that Medication Guides be issued with certain prescribed medications and biological products when the Agency determines that certain information is necessary to prevent serious adverse effects, that patient decision-making should be informed by information about a known serious side effect with a product, or when patient adherence to directions for the use of a product are essential to its effectiveness. During the drug distribution process, if a Medication Guide has been developed for a certain class of medications, then one must be provided with every new prescription and refill of that medication.

Copies of the Medication Guides for psychotropic medications can be accessed on the FDA website at:
<http://www.fda.gov/Drugs/DrugSafety/ucm085729.htm>

Usual Recommended Doses of Common Psychotropic Medications

The attached medication charts are intended to reflect usual doses and brief medication information for commonly used psychotropic medications. The tables contain two columns for maximum recommended doses in children and adolescents – the maximum recommended in the FDA approved product labeling, and the maximum recommended in medical and pharmacological literature sources. The preferred drug list of medications potentially prescribed for foster children is the same as for all other Texas Medicaid recipients.

The tables are intended to serve as a resource for clinicians. The tables are not intended to serve as comprehensive drug information references or a substitute for sound clinical judgment in the care of individual patients. Circumstances may dictate the need for the use of higher doses in specific patients. In these cases, careful documentation of the rationale for the higher dose should occur, and careful monitoring and documentation of response to treatment should be performed. If the use of higher medication doses does not result in improvement in the patient's clinical status within a reasonable time period (e.g., 2-4 weeks), then the dosage should be decreased and other treatment options considered.

Not all medications prescribed by clinicians for psychiatric diagnoses in children and adolescents are included in the following tables. However, in general, medications not listed do not have adequate efficacy and safety information available to support a usual maximum dose recommendation.

See Psychotropic Medication Tables beginning on page 8.

Criteria Indicating Need for Further Review of a Child's Clinical Status

The following situations indicate a need for review of a patient's clinical care. These parameters do not necessarily indicate that treatment is inappropriate, but they do indicate a need for further review.

For a child being prescribed a psychotropic medication, any of the following suggests the need for additional review of a patient's clinical status:

1. Absence of a thorough assessment for the DSM-5 diagnosis(es) in the child's medical record
 2. Four (4) or more psychotropic medications prescribed concomitantly (side effect medications are not included in this count)
 3. Prescribing of:
 - Two (2) or more concomitant stimulants *
 - Two (2) or more concomitant alpha agonists *
 - Two (2) or more concomitant antidepressants
 - Two (2) or more concomitant antipsychotics
 - Three (3) or more concomitant mood stabilizers
- * The prescription of a long-acting and an immediate-release stimulant or alpha agonist of the same chemical entity does not constitute concomitant prescribing.
- Note: When switching psychotropics, medication overlaps and cross taper should occur in a timely fashion, generally within 4 weeks.
4. The prescribed psychotropic medication is not consistent with appropriate care for the patient's diagnosed mental disorder or with documented target symptoms usually associated with a therapeutic response to the medication prescribed.
 5. Psychotropic polypharmacy (2 or more medications) for a given mental disorder is prescribed before utilizing psychotropic monotherapy
 6. The psychotropic medication dose exceeds usual recommended doses (literature based maximum dosages in these tables).
 7. Psychotropic medications are prescribed for children of very young age, including children receiving the following medications with an age of:
 - Stimulants: Less than three (3) years of age
 - Alpha Agonists: Less than four (4) years of age
 - Antidepressants: Less than four (4) years of age
 - Mood Stabilizers: Less than four (4) years of age
 - Antipsychotics: Less than five (5) years of age
 8. Prescribing by a primary care provider who has not documented previous specialty training for a diagnosis other than the following (unless recommended by a psychiatrist consultant):
 - Attention Deficit Hyperactive Disorder (ADHD)
 - Uncomplicated anxiety disorders
 - Uncomplicated depression
 9. Antipsychotic medication(s) prescribed continuously without appropriate monitoring of glucose and lipids at least every 6 months.

Stimulants for treatment of ADHD

Drug (generic)	Drug (brand)+	Initial Dosage	Literature Based Maximum Dosage	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Patient Monitoring Parameters	Black Box Warning**	Warnings and Precautions
Amphetamine mixed salts*	Adderall®	• Age 3-5 years: 2.5 mg/day	Age 3-5 years: 30 mg/day	Approved for children 3 years and older: 40 mg/day	One to three times daily	Baseline and ongoing: height, weight, heart rate, and blood pressure Baseline: Assessment using a targeted cardiac history of the child and the family, and a physical examination of the child with an EKG and/or a pediatric cardiology consult as indicated	• Abuse potential	<ul style="list-style-type: none"> • Risk of sudden death in those with pre-existing structural cardiac abnormalities or other serious heart problems • Hypertension • Potential for psychiatric adverse events (hallucinations, delusional thinking, mania, aggression, etc.) • Stimulants do not appear to affect ultimate adult height. If mild growth suppression occurs, it is likely reversible upon discontinuation of stimulant • Tics • Decreased appetite and weight • Sleep disturbance
	Evekeo®	• Age ≥ 6 years: 5-10 mg/day						
	Adderall®XR	<ul style="list-style-type: none"> • Age 3-5 years: 5mg/day • Age 6-12 years: 5-10 mg/day • Age ≥13 years: 10 mg/day 	Age ≥ 6 years: >50 kg: 60 mg/day	Approved for children 6 years and older: 30 mg/day	Once daily			
Amphetamine base	Adzenys®XR-ODT (oral disintegrating tablet)	• Age ≥ 6 years: 6.3 mg/day (3.1 mg = 5 mg Adderall®XR)	<ul style="list-style-type: none"> Age 6-12 years: 18.8 mg/day Age 13-17 years: 12.5 mg/day 	<ul style="list-style-type: none"> Approved for children 6 years and older: • Ages 6-12 years: 18.8 mg/day • Ages 13-17 years: 12.5 mg/day 	Once daily	<ul style="list-style-type: none"> • Sudden death and serious cardiovascular events (Only boxed warning for amphetamine salts and dextroamphetamine) 		
	Dyanavel®XR (oral suspension)	• Age ≥6 years: 2.5-5 mg/day (2.5 mg = 4 mg Adderall®XR)	Age ≥6 years: 20 mg/day	Approved for children 6 years and older: 20 mg/day	Once daily			
Dextroamphetamine*	Dexedrine®	<ul style="list-style-type: none"> • Age 3-5 years: 2.5 mg/day • Age ≥ 6 years: 5 mg twice daily 	<ul style="list-style-type: none"> Age 3-5 years: 30 mg/day Age ≥ 6 years: >50 kg: 60 mg/day 	Approved for children 3 years and older: 40 mg/day	Once or twice daily			
	Zenzedi®							
	Procentra® (oral suspension)	<ul style="list-style-type: none"> • Age 3-5 years: 5 mg/day • Age ≥ 6 years: 5 mg/day 	Age ≥ 6 years: >50 kg: 60 mg/day	Age ≥ 6 years: 40 mg/day				
	Dexedrine Spansule®							
Lisdexamfetamine	Vyvanse®	<ul style="list-style-type: none"> • Age 3-5 years: 10 mg/day • Age ≥ 6 years: 30 mg/day 	<ul style="list-style-type: none"> Age 3-5 years: 30 mg/day Age ≥ 6 years: 70 mg/day 	Approved for children 6 years and older: 70 mg/day	Once daily			

(Continued on Page 9)

* Generic available

** See the FDA approved product labeling for each medication for the full black box warnings.

+ XR, extended-release

Stimulants for treatment of ADHD (continued)

Drug (generic)	Drug (brand)+	Initial Dosage	Literature Based Maximum Dosage	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Baseline/Monitoring	Black Box Warning	Warnings and Precautions			
Methylphenidate*	Ritalin®	• Age 3-5 years: 2.5 mg twice daily	Age 3-5 years: 20 mg/day Age ≥ 6 years: • ≤50 kg: 60 mg/day • >50 kg: 100 mg/day	Approved for children 6 years and older: 60 mg/day	One to three times daily	See above	See above	See above			
	Methylin® (chewable and oral suspension)	• Age ≥ 6 years: 5 mg twice daily									
	Ritalin®SR	Age ≥ 3 years: 10 mg/day			Approved for children 6 years and older: 60 mg/day				Once daily		
	Methylin®ER										
	Metadate®ER										
	Ritalin®LA	• Age 3-5 years: 10 mg/day • Age ≥ 6 years: 10-20 mg/day			Once daily				See above	See above	See above
	Metadate®CD										
	Quillivant®XR (oral suspension)										
	QuilliChew®ER (chewable)										
	Aptensio®XR										
Concerta®	Age ≥ 3 years: 18 mg/day	Age 3-5 years: 36 mg Age ≥ 6 years: 108 mg/day	Approved for children 6 years and older: • Age 6-12 years: 54 mg/day • Age 13-17 years: lesser of 72 mg/day or 2 mg/kg/day, whichever is less	Once daily							
Daytrana®TD patch †	Age ≥ 3 years: 10 mg/day	Age 3-5 years: 20 mg Age ≥ 6 years: 30 mg/day	Approved for children 6 years and older: 30 mg/day	Once daily							
Dexmethylphenidate*	Focalin®	• Age 3-5 years: 2.5mg/day • Age ≥ 6 years: 2.5 mg twice daily	Age 3-5 years: 10 mg/day	Approved for children 6 years and older: 20 mg/day	Twice daily						
	Focalin®XR	• Age 3-5 years: 5 mg/day • Age ≥ 6 years: 5-10 mg/day	Age ≥ 6 years: 50 mg/day	Approved for children 6 years and older: 30 mg/day	Once daily						

* Generic available

** See the FDA approved product labeling for each medication for the full black box warnings.

† IR, immediate release; SR, sustained-release formulation; CD, combined immediate release and extended release; ER and XR, extended-release; LA, long-acting; TD, transdermal

‡ Daytrana®TD patch: Post marketing reports of acquired skin depigmentation or hypopigmentation of the skin

Other ADHD Treatments

Drug (generic)	Drug (brand)+	Initial Dosage	Literature Based Maximum Dosage	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Baseline/Monitoring	Black Box Warning	Warnings and Precautions
Atomoxetine	Strattera®	<ul style="list-style-type: none"> Age ≥ 6 years and weight ≤70 kg: 0.5 mg/kg/day Age ≥ 6 years and weight >70 kg: 40 mg/day 	Age ≥ 6 years: 1.8 mg/kg/day or 100 mg/day, whichever is less	Approved for treatment of ADHD (age 6-17 years): 1.4 mg/kg/day or 100 mg/day, whichever is less	Once or twice daily	<ul style="list-style-type: none"> Baseline and ongoing: height, weight, heart rate, and blood pressure Onset of therapeutic effect typically delayed 3 weeks 	Suicidal ideation in children and adolescents being treated for ADHD	<ul style="list-style-type: none"> Severe liver injury Contraindicated to use within 14 days of an MAOI Increased blood pressure and heart rate Psychiatric adverse events Priapism (rare)
Clonidine*	Catapres® (IR)	<ul style="list-style-type: none"> Age ≥ 6 years and weight <45 kg: 0.05 mg/day Age ≥ 6 years and weight >45 kg: 0.1 mg/day 	Age ≥ 6 years AND <ul style="list-style-type: none"> Weight 27-40.5 kg: 0.2 mg/day Weight 40.5-45 kg: 0.3 mg/day Weight >45 kg: 0.4 mg/day 	Not approved for treatment of ADHD in children and adolescents	One to four times daily	<ul style="list-style-type: none"> Baseline and ongoing: heart rate and blood pressure Personal and family cardiovascular history 	None	<ul style="list-style-type: none"> Hypotension Bradycardia Syncope Sedation/Somnolence Taper, do not discontinue abruptly
	Kapvay® (ER)	Age ≥ 6 years: 0.1 mg/day	Age ≥ 6 years: 0.4 mg/day	Approved for monotherapy and adjunctive therapy to stimulants for treatment of ADHD (age 6-17 years): 0.4 mg/day	Once or twice daily			
Guanfacine*	Tenex® (IR)	<ul style="list-style-type: none"> Age ≥ 6 years and weight <45 kg: 0.5 mg/day Age ≥ 6 years and weight > 45 kg: 1 mg/day 	Age ≥ 6 years AND <ul style="list-style-type: none"> Weight 27-40.5 kg: 2 mg/day Weight 40.5-45 kg: 3 mg/day Weight >45 kg: 4 mg/day 	Not approved for children and adolescents	One to four times daily	<ul style="list-style-type: none"> Baseline and ongoing: heart rate and blood pressure Personal and family cardiovascular history 	None	<p>CAUTION IF USED WITH ANTIPSYCHOTICS (↓ BP)</p>
	Intuniv® (ER)	Age ≥ 6 years: 1 mg/day	<ul style="list-style-type: none"> Age 6-12 years: 4 mg/day Age 13-17 years: 7 mg/day 	Approved for monotherapy and adjunctive therapy to stimulants for treatment of ADHD <ul style="list-style-type: none"> Age 6-12 years: 4 mg/day Age 13-17 years: 7 mg/day 	Once daily			
Bupropion*	Wellbutrin®	Age ≥ 6 years: 3 mg/kg/day or 150 mg/day, whichever is less	Age ≥ 6 years: 6 mg/kg/day or 300 mg/day with no single dose >150 mg, whichever is less	Not approved for children and adolescents	One to three times daily	<ul style="list-style-type: none"> Blood pressure and Pulse Mental status exam and suicide assessment 	Increased risk of suicidal thinking and behavior (suicidality) in short-term studies in children and adolescents with major depressive disorder (MDD) and other psychiatric disorders	<ul style="list-style-type: none"> Lowers seizure threshold (use caution with other agents that may lower seizure threshold-e.g. antipsychotics, TCAs, excessive alcohol) Discontinuation syndrome Activation of mania/ hypomania Suicidal ideation Contraindicated for use within 14 days of an MAOI
	Wellbutrin®SR	Same as above	400 mg/day		Once or twice daily			
	Wellbutrin®XL	Same as above	450 mg/day		Once daily			
Imipramine*	Tofranil®	Age ≥ 6 years: 1 mg/kg/day or 25 mg/day, whichever is less	Age ≥ 6 years: 4 mg/kg/day or 200 mg/day, whichever is less	Approved for treatment of enuresis in children <ul style="list-style-type: none"> Age 6-11 years: 2.5 mg/kg/day or 50 mg/day, whichever is less Age ≥ 12 years: 2.5 mg/kg/day or 75 mg/day, whichever is less Approved for treatment of depression ≥ 12 years: 100 mg/day	Twice daily	<ul style="list-style-type: none"> CBC Blood pressure and Pulse EKG Mental status exam and suicide assessment 	<ul style="list-style-type: none"> Caution with cardiac disease Cardiac conduction abnormalities Orthostatic hypotension Activation of mania/ hypomania Anticholinergic and cognitive adverse effects Lowers seizure threshold Discontinuation syndrome 	
Nortriptyline*	Aventyl®	Age ≥ 6 years: 0.5 mg/kg/day	Age ≥ 6 years: 2 mg/kg/day or 100 mg/day, whichever is less	Not approved for children and adolescents	Twice daily	<ul style="list-style-type: none"> CBC Blood pressure and Pulse EKG Mental status exam and suicide assessment 		<ul style="list-style-type: none"> Contraindicated for use within 14 days of an MAOI Use caution in those with history of suicide attempts; may be cardiotoxic in overdose
	Pamelor®							
	Nortrilen®							

* Generic available

+ IR, immediate release; SR, sustained-release formulation; ER, extended-release; XL, extended-length

Antidepressants, SSRIs

Drug (generic)	Drug (brand)+	Initial Dosage	Literature Based Maximum Dosage	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Patient Monitoring Parameters	Black Box Warning**	Warnings and Precautions
Citalopram*	Celexa®	<ul style="list-style-type: none"> Age 6-11 years: 10 mg/day Age ≥ 12 years: 20 mg/day 	<ul style="list-style-type: none"> Age ≥ 6 years: 40 mg/day 	Not approved for children and adolescents	Once daily	<ul style="list-style-type: none"> Pregnancy test – as clinically indicated Monitor for emergence of suicidal ideation or behavior Monitor weight and growth Obtain serum sodium if symptoms of hyponatremia occur (e.g. headaches, confusion, etc.) 	Increased risk compared to placebo of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults in short-term studies of major depressive disorder (MDD) and other psychiatric disorders	<ul style="list-style-type: none"> Suicidal ideation Activation of mania/hypomania QTc prolongation potential (citalopram, fluoxetine, possibly escitalopram) Discontinuation syndrome Abnormal bleeding Contraindicated to use within 14 days of an MAOI; for fluoxetine, do not start MAOI for 5 weeks after fluoxetine discontinuation Serotonin Syndrome Hyponatremia risk
Escitalopram*	Lexapro®	<ul style="list-style-type: none"> Age 6-11 years: 5 mg/day Age ≥ 12 years (MDD): 10 mg/day 	<ul style="list-style-type: none"> Age 6-11 years: 20mg/day Age ≥ 12 years: 30 mg/day 	<ul style="list-style-type: none"> Not approved for children Approved for treatment of MDD in adolescents (age 12-17 years): 20 mg/day 				
Fluoxetine*	Prozac®	<ul style="list-style-type: none"> Age 6-11 years: 5-10 mg/day Age ≥ 12 years: 10 mg/day 	<ul style="list-style-type: none"> Age ≥ 6 years: 60/day 	<ul style="list-style-type: none"> Approved for treatment of MDD (age 8-18 years): 20 mg/day Approved for treatment of OCD (age 7-17 years): 60 mg/day 				
Paroxetine*	Paxil®	<ul style="list-style-type: none"> Children: Not recommended Age ≥ 12 years: 10 mg 	<ul style="list-style-type: none"> Children: Not recommended Age ≥ 12 years: 40 mg 	Not approved for children and adolescents				
	Paxil®CR	<ul style="list-style-type: none"> Children: Not recommended Age ≥ 12 years: 25 mg 	<ul style="list-style-type: none"> Children: Not recommended Age ≥ 12 years: 50 mg 					
Fluvoxamine*	Luvox®	Age ≥ 8 years: 25 mg/day	Age 8-11 years: 200 mg/day	Approved for treatment of OCD (age 8-17 years): <ul style="list-style-type: none"> Ages 8-11 years: 200 mg/day Ages 12-17 years: 300 mg/day 	Daily doses >50 mg should be divided			
	Luvox®CR	Age ≥ 8 years: 100 mg/day	Age 12-17 years: 300 mg/day					
Sertraline*	Zoloft®	<ul style="list-style-type: none"> Age 6-12 years: 12.5-25 mg/day Age 13-17 years: 25-50 mg/day 	<ul style="list-style-type: none"> Age ≥ 6 years: 200 mg/day 	Approved for treatment of OCD (age 6-17 years): 200 mg/day	Once daily			
Vilazodone	Viibryd®	Insufficient Evidence	Insufficient Evidence	Not approved for children and adolescents	Insufficient Evidence			

* Generic available

+ CR, controlled-release

**** From Boxed Warning in FDA approved labeling for Antidepressants (SSRIs, SNRIs and Other Mechanisms):** Patients of all ages who are started on antidepressant therapy should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior. Families and caregivers should be advised of the need for close observation and communication with the prescriber. Both patients and families should be encouraged to contact the clinician if depression worsens, the patient demonstrates suicidal behavior or verbalizations, or if medication side effects occur. The appropriate utilization of non-physician clinical personnel who are knowledgeable of the patient population can aid in increasing the frequency of contact between the clinic and the patient/parent.

Antidepressants, SNRIs

Drug (generic)	Drug (brand)+	Initial Dosage	Literature Based Maximum Dosage	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Patient Monitoring Parameters	Black Box Warning	Warnings and Precautions
Venlafaxine*	Effexor®	Age 7-17 years: 37.5 mg/day	<ul style="list-style-type: none"> Age 7-11 years: 150 mg/day Age 12-17 years: 375 mg/day 	Not approved for children and adolescents	IR: Two to three times daily	<ul style="list-style-type: none"> Pregnancy test – as clinically indicated 	<p>Increased risk compared to placebo of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults in short-term studies of major depressive disorder (MDD) and other psychiatric disorders</p>	<ul style="list-style-type: none"> Suicidal ideation Abnormal bleeding Severe skin reactions Discontinuation syndrome Activation of mania/hypomania Hepatotoxicity Elevated blood pressure and pulse Serotonin Syndrome Seizures Hyponatremia Contraindicated for use within 14 days of an MAOI
	Effexor®XR				XR: Once daily			
Duloxetine	Cymbalta®	Age 7-17 years: 30 mg/day	Age 7-17 years: 120 mg/day	Approved for treatment of Generalized Anxiety Disorder Age 7-17 years: 120 mg/day	Once or twice daily	<ul style="list-style-type: none"> Monitor for emergence of suicidal ideation or behavior 		
Desvenlafaxine	Pristiq®	<ul style="list-style-type: none"> Children: Insufficient Evidence Age 12-17 years: 50 mg/day 	<ul style="list-style-type: none"> Children: Insufficient Evidence Age 12-17 years: 100 mg/day 	Not approved for children and adolescents	Once daily	<ul style="list-style-type: none"> Blood pressure during dosage titration and as clinically indicated Monitor weight and growth Hepatic function testing – baseline and as clinically indicated 		
Levomilnacipram	Fetzima®	Insufficient Evidence	Insufficient Evidence	Not approved for children and adolescents	Insufficient Evidence	<ul style="list-style-type: none"> CBC and EKG at baseline and as clinically indicated for Clomipramine 		
Clomipramine*	Anafranil®	Age 10-17 years: 25 mg/day	Age 10-17 years: 3 mg/kg/day or 200 mg/day, whichever is less	Approved for treatment of OCD: Age 10-17 years: 3 mg/kg/day or 200 mg/day, whichever is less	Once daily			

Antidepressants, Other Mechanisms

Drug (generic)	Drug (brand)+	Initial Dosage	Literature Based Maximum Dosage	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Patient Monitoring Parameters	Black Box Warning	Warnings and Precautions
Mirtazapine*	Remeron®	Age ≥ 3 years: 7.5 mg/day	Age ≥ 3 years: 45 mg/day	Not approved for children and adolescents	Once daily	<ul style="list-style-type: none"> Pregnancy test – as clinically indicated Monitor for emergence of suicidal ideation or behavior Blood pressure during dosage titration and as clinically indicated 	<p>Increased the risk compared to placebo of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults in short term studies of major depressive disorder (MDD) and other psychiatric disorders</p>	<ul style="list-style-type: none"> Suicidal ideation Abnormal bleeding Weight gain Discontinuation syndrome Activation of mania/hypomania Orthostatic hypotension and syncope Serotonin Syndrome Hyponatremia Contraindicated for use within 14 days of an MAOI
Vortioxetine	Trintellix®	Insufficient Evidence	Insufficient Evidence	Not approved for children and adolescents	Insufficient Evidence	<ul style="list-style-type: none"> Monitor weight and height Serum cholesterol levels CBC baseline and periodically 		

* Generic Available

+ XR, extended-release

Antipsychotics: Second Generation (Atypical)

Drug (generic)	Drug (brand)+	Initial Dosage	Literature Based Maximum Dosage	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Patient Monitoring Parameters	Black Box Warning	Warnings and Precautions
Aripiprazole*	Abilify® Abilify Discmelt® (oral disintegrating tab) Abilify® (oral solution)	Age ≥ 4 years: 2 mg/day	• Age 4-11 years: 15 mg/day • Age ≥12 years: 30 mg/day	• Approved for treatment of Bipolar Mania or Mixed Episodes (age 10-17 years) and Schizophrenia (13-17 years): 30 mg/day • Approved for treatment of irritability associated with Autistic Disorder (age 6-17 years): 15 mg/day	Once daily	• Fasting plasma glucose level or hemoglobin A1c – at baseline, at 3 months, then every 6 months. • Lipid screening –at baseline, at 3 months, then every 6months.	Increased the risk of suicidal thoughts and behavior in short-term studies in children, adolescents, and young adults with major depressive disorder and other psychiatric disorders	
Quetiapine*	Seroquel® Seroquel®XR (brand only)	• Age 5- 9 years: 12.5-25 mg/day • Age 10-17 years: 50 mg/day	• Age 5- 9 years: 400mg/day • Age 10-17 years: 800 mg/day	• Approved for treatment of Bipolar Mania (age 10-17 years): 600 mg/day • Approved for treatment of Schizophrenia (13-17 years): 800 mg/day	IR: One to three times daily XR: Once daily	• CBC as clinically indicated. • Pregnancy test – as clinically indicated		
Olanzapine*	Zyprexa® Zyprexa Zydis®	• Age 4-5 years: 1.25 mg/day • Age 6-12years: 2.5 mg/day • Age ≥ 13years: 2.5-5 mg/day	• Age 4-5 years: 12.5 mg/day • Age 6-17 years: 20 mg/day	Approved for treatment of Bipolar Mania or Mixed Episodes and Schizophrenia (age 13- 17 years): 20 mg/day	Once daily	• Blood pressure, pulse rate, height, weight and BMI measurement – at every visit • Sexual function– inquire for evidence of galactorrhea/ gynecostasia, menstrual disturbance, libido disturbance or erectile/ ejaculatory disturbances in males (Priapism has been reported with loiperidone, Risperidone and Ziprasidone). This inquiry should be done at each visit for the first 12 months and every 6 months thereafter.	None related to youth	• Extrapyramidal side effects
Risperidone*	Risperdal® Risperdal M-Tab® (oral disintegrating tab) Risperdal® (oral solution)	• Age 4-5 years: ○ <20 kg: 0.25 mg/day ○ >20 kg: 0.5 mg/day • Age ≥6 years: 0.5 mg/day	• Age 4-11 years: 3 mg/day • Age ≥12 years: 6 mg/day	• Approved for treatment of Schizophrenia (age 13-17 years) and Bipolar Mania or Mixed Episodes (age 10-17 years): 6mg/day • Approved for treatment of irritability associated with autistic disorder (age 5-16 years): 3 mg/day	Once or twice daily		None related to youth	• Neuroleptic Malignant Syndrome • Tardive Dyskinesia • Hyperglycemia and Diabetes Mellitus • Prolactinemia and gynecostasia (most common with risperidone and paliperidone)
Clozapine*	Clozaril® FazaClo® (oral disintegrating tablet) Versacloz® oral suspension	• Age 8-11 years: 6.25-12.5 mg/day • Age ≥ 12 years: 6.25-25 mg/day	• Age 8-11 years: 150-300 mg/day • Age ≥ 12 years: 600 mg/day Target serum clozapine level of 350 ng/mL for optimal efficacy	Not approved for children and adolescents	Once or twice daily	• EPS evaluation (examination for rigidity, tremor, akathisia) – before initiation of any antipsychotic medication, then weekly for the first 2 weeks after initiating treatment with a new antipsychotic or until the dose has been stabilized and weekly for 2 weeks after a dose increase. • Tardive Dyskinesia evaluation – every 3 months. .	• Risk of life threatening agranulocytosis • Seizures • Myocarditis • Other adverse cardiovascular and respiratory effects	• Weight gain • Dyslipidemia • Orthostatic Hypotension
Asenapine	Saphris® (sublingual tablet)	• Age ≥ 10 years: 2.5 mg twice daily	Age ≥ 10 years: 10 mg twice daily	Approved for acute treatment of Bipolar Mania and Mixed Episodes (age 10-17 years): 10 mg twice daily	Twice daily. Avoid eating or drinking for 10 minutes after sublingual administration		None related to youth	• Leukopenia, neutropenia, and agranulocytosis • Lowers seizure threshold
Iliperidone**	Fanapt®	Insufficient Evidence	Insufficient Evidence	Not approved for children and adolescents	Insufficient Evidence		None related to youth	• Cognitive and motor impairment potential
Paliperidone*	Invega®	• Children: Insufficient Evidence • Adolescents: (Age ≥ 12 years): 3 mg/day	• Children: Insufficient Evidence • Adolescents (Age ≥ 12 years), Schizophrenia: ○ Weight < 51 kg: 6 mg/day ○ Weight ≥ 51 kg: 12 mg/day	Approved for treatment of Schizophrenia (age 12-17 years): • Weight < 51 kg: 6 mg/day • Weight ≥ 51 kg: 12 mg/day	Once daily	• Vision questionnaire – ask whether the patient has experienced a change in vision and should specifically ask about distance vision and blurry vision-yearly. • EKG - Baseline and as clinically indicated	None related to youth	• Hyperthermia • Dysphagia • Extrapyramidal side effects
Ziprasidone*	Geodon®	• Bipolar Disorder (age 10-17 years): 20 mg/day • Tourette's Disorder: 5 mg/day	• Bipolar Disorder (age 10-17 years) ○ Weight ≤ 45 kg: 80 mg/day ○ Weight > 45 kg: 160 mg/day • Tourette's Disorder: 40 mg/day	Not approved for children and adolescents	Twice daily; take with ≥500 calorie meal	• Clozapine Monitoring Parameters: Clozapine is associated with severe neutropenia (absolute neutrophil count (ANC) less than 500/μL). The requirements to prescribe, dispense, and receive clozapine are incorporated into a single, shared program called the Clozapine Risk Evaluation and Mitigation Strategy (REMS).	None related to youth	• Olanzapine can cause a rare but serious skin reaction known as DRESS (Drug Reaction with Eosinophilia and Systemic Symptoms). Presence of a fever with a rash and swollen lymph glands, or swelling to the face requires immediate medical attention.
Lurasidone	Latuda®	Insufficient Evidence	Insufficient Evidence	Not approved for children and adolescents	Insufficient Evidence Once daily taken with >350 calorie meal		None related to youth	
Brexiprazole	Rexulti®	Insufficient Evidence	Insufficient Evidence	Not approved for children and adolescents	Insufficient Evidence	• Prescribers and pharmacies must certify the use of Clozapine at www.clozapinerems.com .	Antidepressants increased the risk of suicidal thoughts and behavior in children, adolescents, and young adults in short-term studies.	

* Generic available

+ XR, extended-release

** While iliperidone alone can cause QTc prolongation, concomitant administration with a CYP2D6 inhibitor (e.g., paroxetine) or a CYP3A4 inhibitor (e.g., ketoconazole) can double QTc prolongation compared with administering iliperidone alone.

No long-acting injectable antipsychotic formulations are FDA-approved for use in children and adolescents

Antipsychotics: First Generation (Typical)

Drug (generic)	Drug (brand)	Initial Dosage	Literature Based Maximum Dosage	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Patient Monitoring Parameters	Black Box Warning	Warnings and Precautions
Chlorpromazine*	Thorazine®	<ul style="list-style-type: none"> Age > 6 months: 0.25 mg/lb every 4-6 hours, as needed Adolescents: 10-25 mg/dose every 4-6 hours 	<ul style="list-style-type: none"> Age < 5 years: 40 mg/day Age 5-12 years: 75 mg/day Age > 12 years: 800 mg/day 	<p>Approved for treatment of severe behavioral problems (age 6 months-12 years)</p> <ul style="list-style-type: none"> Outpatient Children: 0.55 mg/kg every 4-6 hours, as needed Inpatient Children: 500 mg/day <p>Approved for the management of manifestations of Psychotic Disorders (age > 12 years): 1000 mg/day</p>	One to six times daily	Same as Second Generation Antipsychotics	None related to youth	<ul style="list-style-type: none"> Tardive Dyskinesia Neuroleptic Malignant Syndrome Leukopenia, neutropenia, and agranulocytosis Drowsiness
Haloperidol*	Haldol®	<ul style="list-style-type: none"> Age 3-12 years weighing <ul style="list-style-type: none"> ○ 15-40 kg: 0.025-0.05 mg/kg/day ○ ≥ 40 kg: 1 mg/day Age > 12 : 1 mg/day 	<ul style="list-style-type: none"> Age 3-12 years: 0.15 mg/kg/day or 6 mg/day, whichever is less Age >12 years <ul style="list-style-type: none"> ○ Acute agitation: 10 mg/dose ○ Psychosis: 15 mg/day ○ Tourette's Disorder: 15 mg/day 	<p>Approved for treatment of Psychotic Disorders, Tourette's Disorder, and severe behavioral problems (age ≥3 years):</p> <ul style="list-style-type: none"> Psychosis: 0.15 mg/kg/day Tourette's Disorder and severe behavioral problems: 0.075 mg/kg/day Severely disturbed children: 6 mg/day 	One to three times daily		None related to youth	<ul style="list-style-type: none"> Orthostatic hypotension EKG changes Extrapyramidal symptoms Ocular changes Hyperprolactinemia
Perphenazine*	Trilafon®	<ul style="list-style-type: none"> Age 6-12 years: Insufficient Evidence Age > 12 years: 4-16 mg two to four times daily 	<ul style="list-style-type: none"> Age 6-12 years: Insufficient Evidence Age > 12 years: 64 mg/day 	<p>Approved for treatment of psychotic disorders (age ≥12 years):</p> <ul style="list-style-type: none"> Outpatient: 24 mg/day Inpatient: 64 mg/day 	Two to four times daily		None related to youth	<ul style="list-style-type: none"> Anticholinergic effects (constipation, dry mouth, blurred vision, urinary retention)
Pimozide	Orap®	Age ≥7 years: 0.05 mg/kg	<ul style="list-style-type: none"> Age 7-12 years: 6 mg/day or 0.2 mg/kg/day, whichever is less Age ≥ 12 years: 10 mg/day or 0.2 mg/kg/day, whichever is less 	<p>Approved for treatment of Tourette's Disorder (age ≥12 years):</p> <p>10 mg/day or 0.2 mg/kg/day, whichever is less</p>	Once or twice daily		None	<ul style="list-style-type: none"> Risk of prolonged QTc interval and torsades de pointes (particularly with pimozide)

* Generic available

Mood Stabilizers

Drug (generic)	Drug (brand)+	Initial Dosage	Target Dosage Range	Literature Based Maximum Dosage	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Patient Monitoring Parameters	Black Box Warning	Warnings and Precautions
Carbamazepine*	Epitol® (tab) Tegretol® (tab, oral suspension, chewable)	• Age 4-5 years: 10-20 mg/kg/day	• Age 4-5 years: 35 mg/kg/day	• Age 4-5 years: 35 mg/kg/day	Approved for treatment of Seizure Disorders in all ages	Two to four times daily	<ul style="list-style-type: none"> CBC with differential --- baseline and 1 to 2 weeks after each dose increase, annually, and as clinically indicated Electrolytes --- baseline and 1 to 2 weeks after each dose increase, annually, and as clinically indicated Hepatic function - baseline, monthly for first three months, annually and as clinically indicated. Pregnancy Test --- baseline as appropriate, and as clinically indicated Carbamazepine levels ---obtain 1 week after initiation and 3-4 weeks after dose adjustment, then as clinically indicated 	• Stevens-Johnson Syndrome	<ul style="list-style-type: none"> Stevens-Johnson Syndrome Aplastic anemia Suicidality Teratogenicity Neutropenia and agranulocytosis Hyponatremia
	Tegretol®XR (tab) Carbatrol® (extended release capsule) Equetro® (extended release capsule)	• Age 6-12 years: 10 mg/kg/day or 200 mg/day	• Ages 6-12 years: 400-800 mg/day	• Ages 6-12 years: 800 mg/day	• Age < 6 years: 35 mg/kg/day	• Age 6-15 years: 1000 mg/day	Twice daily	<ul style="list-style-type: none"> For patients with Asian descent, genetic test for HLA- B*1502 at baseline (prior to the initiation of carbamazepine). May use results of previously completed testing. Patients testing positive for the allele should not use carbamazepine unless benefit outweighs the risk Consider HLA-A*3101 genetic testing at baseline for those to be considered at high risk (most common in Asian, Native American, European, and Latin American descents) Monitor for the emergence of suicidal ideation or behavior <p>Usual therapeutic levels 4-12 mcg/ml</p>	• Aplastic Anemia/granulocytosis
Divalproex Sodium*	Depakote® delayed- release tablets Depakote® ER extended-release tablets Depakote® sprinkles	Age ≥6 years: 10-15 mg/kg/day	Age ≥6 years: 30-60 mg/kg/day	Age ≥6 years: Serum level: 125 µg/mL, or 60 mg/kg/day	Approved for treatment of Seizure Disorders (age ≥ 10 years) Maximum dose based upon serum level: 50-100 µg/mL, or 60 mg/kg/day	One to three times daily	<ul style="list-style-type: none"> CBC - with differential and platelet count - baseline then 1 to 2 weeks after each dosage increase, every 3 months for the first year of treatment, then annually and as clinically indicated Comprehensive Metabolic Panel (hepatic function, serum creatinine, BUN and electrolytes) – baseline, every 3 months for the first year of treatment, then annually and as clinically indicated. Pregnancy Test – baseline as appropriate, and as clinically indicated Valproic acid level – 1-2 weeks after initiation and dosage change, then as clinically indicated. Weight – baseline, quarterly for the first year of treatment, then annually and as clinically indicated Monitor for the emergence of suicidal ideation or behavior Usual therapeutic trough levels for bipolar disorder is 50- 125 mcg/ml for Valproic acid and Divalproex delayed release (Depakote®). <p>For divalproex extended release (Depakote® ER) it is 85 – 125 mcg/ml (trough) for the treatment of acute mania. A lower therapeutic trough level may be needed with Divalproex extended release for maintenance treatment.</p> <p>For extended release products, a trough level is considered to be 18 to 24 hours after the last dose</p>	• Hepatotoxicity • Teratogenicity • Pancreatitis	<ul style="list-style-type: none"> Hepatotoxicity Pancreatitis Urea cycle disorders Teratogenicity Suicidal ideation Neutropenia and leukopenia (significant increased risk with quetiapine co-administration) Thrombocytopenia Hyperammonemia Multi-organ hypersensitivity reaction Withdrawal seizures Polycystic ovarian syndrome Weight gain Alopecia
Lithium*	Eskalith®	• Age 6-11 years: Lesser of 15-20 mg/kg/day or 150mg twice per day • Age ≥ 12 years: Lesser of 15-20 mg/kg/day or 300 mg twice per day	Dose adjustment based upon serum level 12 hour post dose serum level: 0.6-1.2 mEq/L	Age ≥6 years: Serum level: 1.2 mEq/L, or 1800 mg	Approved for treatment of manic episodes and maintenance of Bipolar Disorder (age ≥ 12 years) Maximum dose based upon 12 hour post dose serum level: 1.2 mEq/L	One to four times daily	<ul style="list-style-type: none"> EKG – baseline, yearly and as clinically indicated CBC – baseline, yearly and as clinically indicated Thyroid studies – baseline; then TSH every 6 months and as clinically indicated Comprehensive Metabolic Panel (BUN, creatinine, glucose, calcium, and electrolytes)-baseline, 3 months, annually and as clinically indicated. Caution: BUN:serum creatinine ratio >20 may be an indication of dehydration. UA - baseline and as clinically indicated Pregnancy Test - as clinically indicated Lithium Levels – one week (i.e., 5-7 days) after initiation or dosage change, 3 months after initiation, and as clinically indicated; for maintenance treatment every 6 months, and as clinically indicated Weight – baseline, every 6 months and as clinically indicated Usual trough therapeutic level: 0.6-1.2 meq/L (12 hour post dose) 	Toxicity above therapeutic serum levels	<ul style="list-style-type: none"> Toxicity above therapeutic serum levels Chronic renal function impairment Special risk patients: those with significant renal or cardiovascular disease, severe debilitation, dehydration, or sodium depletion Polyuria Tremor Diarrhea Nausea Hypothyroidism Teratogenicity
	Eskalith®CR								
	Lithobid®(ER)								

(Continued on Page 16)

* Generic Available
+ ER and XR, extended-release; CR, controlled release

Mood Stabilizers (continued)

Drug (generic)	Drug (brand)+	Initial Dosage	Target Dosage Range	Literature Based Maximum Dosage	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Patient Monitoring Parameters	Black Box Warning	Warnings and Precautions
Lamotrigine*	Lamictal®	<ul style="list-style-type: none"> Age 6-11 years: 2.5 mg/day Age ≥12 years: 25 mg/day (increase by 25 mg every 2 weeks) 	<p>Age 6-11 years</p> <ul style="list-style-type: none"> Monotherapy: 4.5-7.5 mg/kg/day With Valproate: 1-3 mg/kg/day With Valproate and EIAEDs †: 1-5 mg/kg/day With EIAEDs †: 5-15 mg/kg/day <p>Age ≥12 yearsw</p> <ul style="list-style-type: none"> Monotherapy: 225-375 mg/day With Valproate: 100-200 mg/day With Valproate and EIAEDs †: 100-400 mg/day With EIAEDs †: 300-500 mg/day 	Age ≥6 years: 15 mg/kg/day or 500 mg/day, whichever is less	<p>Approved for adjunctive therapy for Seizure Disorders:</p> <p>Age 2-12: 400 mg/ day</p> <p>Age >12: 500 mg/day (use > 200mg/day in adults for bipolar depression has not conferred additional efficacy)</p> <p>Safety and effectiveness for treatment of Bipolar Disorder in patients younger than 18 years had not been established</p>	Once or twice daily	<ul style="list-style-type: none"> Renal Function - baseline and as clinically indicated Hepatic Function - baseline and as clinically indicated Pregnancy Test - baseline and as clinically indicated CBC – baseline and as clinically indicated Monitor for the emergence of suicidal ideation or behavior Monitor for rash, especially during the first two months of therapy 	Serious rashes including Stevens-Johnson syndrome	<ul style="list-style-type: none"> Dermatological reactions Potential Stevens- Johnson Syndrome; risk increased with too-rapid titration Multi-organ Hypersensitivity reactions and organ failure Suicidal ideation Aseptic meningitis Concomitant use with Divalproex increases serum Lamotrigine levels significantly (increased risk of rash/SJS without lamotrigine dose adjustment) Concomitant use with enzyme induced AEDs (Carbamazepine, Phenytoin, Phenobarbital, Primidone) reduces serum lamotrigine levels significantly (reduced lamotrigine efficacy possible without lamotrigine dose adjustment) Concomitant use with oral contraceptives increases lamotrigine clearance Withdrawal seizures
Oxcarbazepine*	Trileptal®	8-10 mg/kg/day	<p>Monotherapy (based on weight):</p> <ul style="list-style-type: none"> 20-24.9 kg: 600-900 mg/day 25-34.9 kg: 900-1200 mg/day 35-44.9 kg: 900-1500 mg/day 45-49.9 kg: 1200 – 1500 mg/day 50-59.9 kg: 1200-1800 mg/day 60-69.9 kg: 1200-2100 mg/day ≥70 kg: 1500-2100 mg/day 	<ul style="list-style-type: none"> Age 7-12 years: 60 mg/kg/day or 1500 mg/day Age 13-17 years: 60 mg/kg/day or 2100 mg/day 	<p>Approved for treatment of Seizure Disorders as mono- therapy (age ≥ 4 years), or as adjunctive therapy in (age ≥ 2 years): 60 mg/kg/day or 1800 mg/day</p> <p>Safety and effectiveness for treatment of Bipolar Disorder in patients younger than 18 years had not been established</p>	Twice daily	<ul style="list-style-type: none"> CBC with differential – baseline and 1 to 2 weeks after each dose increase, annually, and as clinically indicated Electrolytes – baseline and 1 to 2 weeks after each dose increase; monthly for the first 3 months, then annually, and as clinically indicated Hepatic function - baseline and annually Pregnancy Test – baseline as appropriate and as clinically indicated For patients with Asian descent, genetic test for HLA- B*1502 at baseline (prior to the initiation of oxcarbazepine). May use results of previously completed testing. Monitor for the emergence of suicidal ideation or behavior Obtain serum sodium if symptoms of hyponatremia occur (headaches, confusion, etc.) 	None	<ul style="list-style-type: none"> Hyponatremia (incidence may be as high as 24% in children) Drug-drug interaction potential Anaphylactic reactions and angioedema Patients with a past history of hypersensitivity reaction to carbamazepine Serious dermatological reactions Withdrawal seizures Cognitive/neuropsychiatric adverse events Multi-organ hypersensitivity Hematologic events

* Generic Available

+ ER and XR, extended-release; CR, controlled release

†EIAED's - Enzyme Inducing Anti-Epileptic Drugs (e.g. Carbamazepine, Phenobarbital, Phenytoin, Primidone)

Sedatives/Hypnotics

Drug (generic)	Drug (brand)	Initial Dosage	Literature Based Maximum Dosage**	FDA Approved Maximum Dosage for Children and Adolescents	Schedule	Black Box Warning**	Warnings and Precautions
Diphenhydramine*	Benadryl®	<ul style="list-style-type: none"> Age 3-5 years: 6.25-12.5 mg (1mg/kg max) Age 5-11 years: 12.5-25 mg Age ≥12 years: 25-50 mg 	<ul style="list-style-type: none"> 25-37 lbs: 12.5 mg 38-49 lbs: 19 mg 50-99 lbs: 25 mg ≥100 lbs: 50 mg Evidence suggests that tolerance develops to the hypnotic effects of diphenhydramine within 5-7 nights of continuous use.	Approved for treatment of insomnia (age ≥12 years): 50 mg at bedtime	Once at bedtime	None	<ul style="list-style-type: none"> Drowsiness Dizziness Dry mouth Nausea Nervousness Blurred vision Diminished mental alertness Paradoxical excitation Respiratory disease Hypersensitivity reactions May lower seizure threshold (avoid in epilepsy)
Trazodone*	Desyrel®	<ul style="list-style-type: none"> Children: Insufficient Evidence Adolescents: 25 mg 	<ul style="list-style-type: none"> Children Insufficient Evidence Adolescents: 100 mg/day 	Not approved for children or adolescents as a hypnotic	Once at bedtime	Increased the risk compared to placebo of suicidal thinking and behavior (Suicidality) in children, adolescents, and young adults in short-term studies of major depressive disorder (MDD) and other psychiatric disorders	<ul style="list-style-type: none"> Serotonin Syndrome Contraindicated for use within 14 days of an MAOI Suicidal ideation Activation of mania/hypomania Discontinuation syndrome Abnormal bleeding QT prolongation and risk of sudden death Orthostatic hypotension and syncope Abnormal bleeding Priapism Hyponatremia Cognitive and motor impairment
Eszopiclone	Lunesta®	Insufficient Evidence	Insufficient Evidence	Not approved for children or adolescents	Once at bedtime	None	<ul style="list-style-type: none"> Complex sleep behaviors possible Abnormal thinking and behavior changes Withdrawal effects Drug abuse and dependence Tolerance
Melatonin	No brand name	<ul style="list-style-type: none"> Age 3-5 years: 0.5mg Age ≥6 years: 1mg 	<ul style="list-style-type: none"> Age 3-5 years: 0.15 mg/kg or 3 mg, whichever is less Age ≥6 years: 0.15mg/kg or 6mg, whichever is less 	Regulated by FDA as a dietary supplement and not as a medication (no FDA approved indications)	Once at bedtime or alternatively, give 5-6 hrs before Dim Light Melatonin Onset (DLMO)	None	<ul style="list-style-type: none"> Sedation May adversely affect gonadal development Should be given directly before onset of sleep is desired due to short half-life
Ramelteon	Rozerem®	Insufficient Evidence	Insufficient Evidence	Not approved for children or adolescents	Insufficient Evidence	None	<ul style="list-style-type: none"> Hypersensitivity reactions Need to evaluate for comorbid diagnoses Abnormal thinking and behavioral changes CNS depression Decreased testosterone Hyperprolactinemia
Hydroxyzine*	Vistaril®	<ul style="list-style-type: none"> Age 3-5 years: 25 mg Age ≥6 years: 50mg 	<ul style="list-style-type: none"> Age 3-5 years: 25 mg Age 6-11 years: 50mg Age 12 years and older: 100 mg 	Approved for treatment of anxiety and tension: <ul style="list-style-type: none"> Age <6 years: 50 mg/day in divided doses Age = 6 years: 50-100 mg/day in divided doses Approved as a sedative when used as a premedication and following general anesthesia: 0.6 mg/kg	Once at bedtime	None	<ul style="list-style-type: none"> Drowsiness Dry mouth Involuntary motor activity Blurred vision, dizziness, diminished mental alertness Paradoxical excitation associated with a small but definite risk of QT interval prolongation and torsades de pointes

* Generic Available

** Maximum doses for the sedative/hypnotics are based upon night time doses to induce sleep in a child with severe insomnia.

Glossary

ANC = ABSOLUTE NEUTROPHIL COUNT

BMI = Body Mass Index. A measure of body fat based upon height and weight.

CBC = Complete blood count. Lab test used to monitor for abnormalities in blood cells, e.g., for anemia.

C_p = Plasma concentration

Serum creatinine = A lab test used to calculate an estimate of kidney function.

EKG = Electrocardiogram

EEG = Electroencephalogram

EPS = Extrapyramidal side effects. These are adverse effects upon movement, including stiffness, tremor, and severe muscle spasm

FDA = U.S. Food and Drug Administration

Hemoglobin A1c = A laboratory measurement of the amount of glucose in the hemoglobin of the red blood cells. Provides a measure of average glucose over the previous 3 months.

LFTs = Liver function tests

MAOIs = Monoamine Oxidase Inhibitors

MRI = Magnetic resonance imaging

PRN = as needed

Prolactin = A hormone produced by the pituitary gland

TFTs = Thyroid Function Tests

References

- Aman MG, Bukstein OG, Gadow KD, et al. What does risperidone add to parent training and stimulant for severe aggression in child attention-deficit/hyperactivity disorder? *J Am Acad Child Adolesc Psych* 2014;53:47-60.
- Biederman et al. A prospective open-label trial of lamotrigine monotherapy in children and adolescents with bipolar disorder. *CNS Neurosci Ther* Apr 2010;16(2):91-102
- Blader JC, Pliszka SR, Kafantaris V, et al. Prevalence and treatment outcomes of persistent negative mood among children with attention-deficit/hyperactivity disorder and aggressive behavior. *J Child Adolesc Psychopharmacol*. 2016;26(2):164-173.
- Bobo WV, Cooper WO, Stein CM, et al. Antipsychotics and the risk of type 2 diabetes mellitus in children and youth. *JAMA Psychiatry* 2013;70:1067-75.
- Bridge JA, Iyengar S, Salary CB, et al. Clinical response and risk for reported suicidal ideal and suicide attempts in pediatric antidepressant treatment. *JAMA* 2007;297:1683-96.
- Correll CU. Monitoring and management of antipsychotic-related metabolic and endocrine adverse events in pediatric patients. *Int Rev Psychiatry* 2008;20(2):195-201.
- Correll CU, Kratochvil CJ, March JS. Developments in pediatric psychopharmacology: focus on stimulants, antidepressants, and antipsychotics. *J Clin Psychiatry* 2011;722:655-70.
- Cooper WO, Habel LA, Sox CM et al. ADHD drugs and serious cardiovascular events in children and young adults. *N Engl J* 2011;365:1896904. Crismon ML, Argo T. The Use of Psychotropic Medication for Children in Foster Care. *Child Welfare* 2009;88:71-100.
- Cutler AJ, Brams M, Bukstein O, et al. Response/remission with guanfacine extended-release and psychostimulants in children and adolescents with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psych* 2014;53:1092-1101.
- De Hert M, Bobbelaere M, Sheridan EM, et al. Metabolic and endocrine adverse effects of second-generation antipsychotics in children and adolescents: A systematic review of randomized, placebo controlled trials and guidelines for clinical practice. *European Psychiatry* 2011;26:144- 58.
- Detke HC, DelBello MP, Landry J, Usher RW. Olanzapine/fluoxetine combination in children and adolescents with bipolar I depression: a randomized, double-blind placebo controlled trial. *J Am Acad Child Adolesc Psych* 2015;54:217-224.
- Dopheide JA, Pliszka SR. Attention-Deficit-Hyperactivity Disorder: An Update. *Pharmacotherapy* 2009;29(6)656-79.
- Emslie GJ, et al. Escalopram in the treatment of adolescent depression: a randomized placebo-controlled multisite trial. *J Am Acad Child Adolesc Psychiatry* 2006;48(7)721-9.
- Etminan M, Carleton B and Brophy JM. Risperidone and Risk of Gynecomastia in Young Men. *J Child Adolesc Psychopharm* 2015;25(9):671- 673.
- Fanton J, Gleason M. Psychopharmacology and preschoolers: a critical review of current conditions. *Child Adolesc Psychiatr Clin N Am* 2009;18(3):753-71.
- Findling RL. Efficacy, long-term safety, and tolerability of ziprasidone in children and adolescents with bipolar disorder. *J Child Adolesc Psychopharm* 2013;23(8):545-57.
- Findling RL, Drury SS, Jensen PS, AACAP Committee on Quality Issues. Practice parameter for the use of atypical antipsychotic medications in children and adolescents. American Academy of Child and Adolescent Psychiatry. Approved by the AACAP Council on August 2, 2011.
- Findling RL, Johnson JL, McClellan J, et al. Double-blind maintenance safety and effectiveness findings from the treatment of early-onset schizophrenia spectrum (TEOSS) study. *J Am Acad Child Adolesc Psychiatry* 2010;49:583-94.

References (continued)

- Findling RL, Reed MD, O’Riordan MA, Demeter CA, Stansbery RJ, McNamara NK. Effectiveness, safety, and pharmacokinetics of quetiapine in aggressive children with conduct disorder. *J Am Acad Child Adolesc Psychiatry* 2006;45:792-800.
- Findling RL, Robb A, McNamara NK, et al. Lithium in the acute treatment of bipolar I disorder: a double-blind placebo-controlled study. *Pediatrics* 2015;13:885-894.
- Gadow KD, Arnold LG, Molina BSG, et al. Risperidone added to parent training and stimulant medication: Effects on attention-deficit/hyperactivity disorder, conduct disorder, and peer aggression. *J Am Acad Child Adolesc Psych* 2014;53:948-959.
- Galling B et al. Type 2 Diabetes Mellitus in Youth Exposed to Antipsychotics: A Systematic Review and Meta-analysis. *JAMA psychiatry*. Online Jan 20, 2016:e1-e13.
- Gandelman K, Alderman JA, Glue P, et al. The impact of calories and fat content of meals on oral ziprasidone absorption: A randomized, open-label, crossover trial. *J Clin Psychiatry* 2009;70:58-62.
- Geller B, Luby JL, Joshi P, et al. A randomized controlled trial of risperidone, lithium, or divalproex sodium for initial treatment of bipolar I disorder, manic or mixed phase, in children and adolescents. *Arch Gen Psych* 2012;69:515-528.
- Gibbons RD, Brown H, Hur K, Davis JM, Mann JJ. Suicidal thoughts and behavior with antidepressant treatment: Reanalysis of the randomized placebo-controlled studies of fluoxetine and venlafaxine. *Arch Gen Psych* 2012;69:580-587.
- Ghanizadeh A et al. A head-to-head comparison of aripiprazole and risperidone for safety and treating autistic disorders. *Child Psychiatry Hum Dev* 2014;45(2):185-192. Geller B, Luby JL, Joshi P, et al. A randomized controlled trial of risperidone, lithium, or divalproex sodium for initial treatment of bipolar I disorder, manic or mixed phase, in children and adolescents (TEAM Study). *Arch Gen Psychiatry* 2012;69:515-28.
- Gleason MM, Egger HL, Emslie GJ, et al. Psychopharmacological treatment for very young children: contexts and guidelines. *J Am Acad Child Adolesc Psychiatry* 2007;46:1532-1572.
- Greenhill L, Kollins S, Abikoff H, et al. Efficacy and safety of immediate-release methylphenidate treatment for preschoolers with ADHD. *J Am Acad Child Adolesc Psychiatry* 2006;45(11):1284–1293.
- Groenman AP, Oosterlaan NN, et al. Stimulant therapy for attention-deficit hyperactivity disorder and risk of developing substance abuse disorder. *Pediatrics* 2014;204:494-501
- Hammond TA, Laughren T, Racoosin J. Suicidality in pediatric patients treated with antidepressant drugs. *Arch Gen Psychiatry*. 2006;63:332- 339.
- Hastard EB et al. ADHD, stimulant treatment, and growth: a longitudinal study. *Pediatrics* 2014;134(4):e935-944.
- Hay W, Levin M, Deterding R, et al. *Current diagnosis and treatment pediatrics*. 20th ed. McGraw-Hill Professional Publishers 2010.
- Hirota T, Schwartz S, Correll CU. Alpha-2 agonists for attention-deficit/hyperactivity disorder in youth: systematic review and meta-analysis of monotherapy and add-on trials to stimulant therapy. *Jour Am Acad Child Adolesc Psych* 2014;53:153-173.
- Jensen KG, Juul K, Fink-Jensen A, Correll CU, Pagsberg AK. Corrected QT changes during antipsychotic treatment of children and adolescents: a systematic review and meta-analysis of clinical trials. *J am Acad Child Adolesc Psych* 2015;54-25-36.
- Johns Hopkins Hospital, Arcara K, Tschudy M. *The Harriet Lane Handbook: A Manual for Pediatric House Officers*. 19th ed. Philadelphia, PA: Mosby/Elsevier 2012.
- Kennard BD, Emslie GJ, Mayes TL, et al. Sequential treatment with fluoxetine and relapse-prevention CBT to improve outcomes in pediatric depression. *Am J Psych* 2014;171:1083-1090.

References (continued)

- Keller MB et al. Efficacy of paroxetine in the treatment of adolescent major depression: a randomized, controlled trial. *J Am Acad Child Adolesc Psychiatry* 2001;40(7):762-72.
- Knapp P, Chait A, Pappadopulos, E, Crystal S, Jensen PS, T-MAY Steering Committee. Treatment of maladaptive aggression in youth. CERT Guidelines I. Engagement, assessment, and management. *Pediatrics* 2012;129(6):e-1562-76.
- Kliegman RM, Stanton B, Geme J, et al. *Nelson textbook of pediatrics*. 19th ed. Saunders Publishers; 2011.
- Kowatch RA, Scheffer RE, Monroe E, et al. Placebo-controlled trial of valproic acid versus risperidone in children 3-7 years of age with bipolar I disorder. *Jour Child Adolesc Psychopharm* 2015;25:306-313.
- Lam RW. Antidepressants and QTc prolongation. *J Psychiatry Neurosci* 2013;38:E5-6.
- Le Noury J et al. Restoring Study 329: efficacy and harms of paroxetine and imipramine in treatment major depression in adolescence. *BMJ* 2015;351:h4320.
- March JS. The preschool ADHD treatment Study (PATS) as the culmination of twenty years of clinical trials in pediatric psychopharmacology. *J Am Acad Child Adolesc Psychiatry* 2011;50(5):427-30.
- McVoy M, Findling RL eds. *Clinical manual of child and adolescent psychopharmacology*, 2nd ed. American Psychiatric Publishing Washington, DC. 2013.
- Miller M, Swanson SA, Azrael D, Pate V, Styrmer T. Antidepressant dose, age, and the risk of deliberate self-harm. *JAMA Intern Med* 2014;174:899-909.
- Nagy P, Nagy A, et al. Functional outcomes from a head-to-head trial of lisdexamfetamine dimesylate and atomoxetine in children and adolescents with attention-deficit/hyperactivity disorder and an inadequate response to methylphenidate. *Eur Child Adolesc Psychiatry*; online May 22, 2015.
- Lelayo R, Yuen K. Pediatric sleep pharmacology. *Child Adolesc Psychiatr Clin N Am* 2012;21:861-83.
- Perrin JM, Friedman RA, Knilans TK, et al. Cardiovascular monitoring and stimulant drugs for Attention Deficit/Hyperactivity Disorder. *Pediatrics* 2008;122:451-453.
- Peukens J. The effects of novel and newly approved antipsychotics on serum prolactin levels: a comprehensive review. *CNS Drugs* 2014;28(5):421-53.
- Pliszka SR, AACAP Work Group on Quality Issues. Practice parameter for the assessment and treatment of children and adolescents with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2007;46:894-921.
- Rahman A et al. Evaluating the incidence of leukopenia and neutropenic with valproate, quetiapine, or the combination in children and adolescents. *Ann Pharmacother* 2009;43(5):822-30.
- Rosato NS, Correll CU, Pappadopulos, E, Chait A, Crystal S, Jensen PS, T-MAY Steering Committee. Treatment of maladaptive aggression in youth. CERT guidelines II. Treatments and ongoing management. *Pediatrics* 2012;129(6):e1577-86.
- Scahill L, Oesterheld, JR. Martin A. Pediatric psychopharmacology II. General principles, specific drug treatments, and clinical practice. In: Lewis M (ed.). *Child and adolescent psychiatry: A comprehensive textbook*. Lippincott Williams & Wilkins, Philadelphia 2007 754-788.
- Scheeringa MS, Weems CF, Cohen JA, et al. Trauma-focused cognitive-behavioral therapy for posttraumatic stress disorder in 3 through 6 year- old children: a randomized clinical trial. *J Child Psychol Psychiatry* 2011;52:853-860.
- Schelleman H, Bilker WB, Strom BL, et al. Cardiovascular events and death in children exposed and unexposed to ADHD agents. *Pediatrics* 2011;127(6):1102-1110.

References (continued)

- Seida JC, Schouten JR, Boylan K, et al. Antipsychotics for children and young adults: a comparative effectiveness review. *Pediatrics* 2012;129:e771-e784.
- Sikich L, Frazier JA, McCelessan J, et al. Double-Blind Comparison of First- and Second-Generation Antipsychotics in Early-Onset Schizophrenia and Schizoaffective Disorder: Findings from the Treatment of Early-Onset Schizophrenia Spectrum Disorders (TEOSS) Study. *Am J Psychiatry* 2008;165(11): 1420-31.
- Southammakosane C, Schmitz, K. Pediatric Psychopharmacology for Treatment of ADHD, Depression, and Anxiety. *Pediatrics* 2015; 136(2):351-59.
- Sporn AL et al. Clozapine treatment of childhood-onset schizophrenia: evaluation of effectiveness, adverse effects, and long-term outcome. *J Am Acad Child Adolesc Psychiatry* 2007;46 (10):1349-56.
- Stigler KA, et al. Paliperidone for irritability in adolescents and young adults with autistic disorder. *Psychopharmacology (Berl)* 2012;223(2)237- 45.
- Subcommittee on Attention-Deficit Hyperactivity Disorder, Steering Committee on Quality Improvement and Management. ADHD: Clinical Practice Guideline for the Diagnosis, Evaluation, and Treatment of Attention-Deficit/Hyperactivity Disorder in Children and Adolescents. *Pediatrics* 2011;128:1007-22.
- van Geijlswijk IM, van der Heijden KB, Egberts ACG, et al. Dose finding of melatonin for chronic idiopathic childhood sleep onset insomnia: an RCT. *Psychopharmacology* 2010;212:379-391.
- Varigonada AL, Jakubovski E, Taylor MJ, et al. Systematic review and meta-analysis: early treatment responses of selective serotonin reuptake inhibitors in pediatric major depressive disorder. *J Am Acad Child Adolesc Psych* 2015;54:557-564.
- Vaughan B, Kratochvil CJ. Pharmacotherapy of pediatric attention-deficit/hyperactivity disorder. *Child Adolesc Psychiatric Clin N Am* 2012;21:941-55.
- Vetter VL et al. Cardiovascular monitoring of children and adolescents receiving medications for ADHD-Scientific Statement. *American Heart Assoc Circulation* 2008;117:2407-2423.
- Wagner KD. A double blind, randomized placebo-controlled trial of escitalopram in the treatment of pediatric depression. *J Am Acad Child Adolesc Psychiatry* 2006;45(3):280-288.
- Wagner KD, Pliszka SR. Treatment of child and adolescent disorders. In: Schatzberg AF, Nemeroff CB (eds). *Textbook of psychopharmacology*, 4th. Ed. American Psychiatric Publishing, Washington, DC, 2009;1309-1371.
- Walkup J, Work Group on Quality Issues. Practice parameter on the use of psychotropic medication in children and adolescents. *J Am Acad Child Adolesc Psychiatry* 2009;48:961-73.
- Wilens TE, Robertson B, Sikirca V, et al. A randomized, placebo-controlled trial of guanfacine extended release in adolescents with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2015;54:916-25.
- Wozniak J, Mick E, Waxmonsky J, et al. Comparison of Open-Label, 8-Week Trials of Olanzapine Monotherapy and Topiramate Augmentation of Olanzapine for the Treatment of Pediatric Bipolar Disorder. *J Child Adolesc Psychopharmacol* 2009;19(5):539-45.
- Zolaly MA. Histamine H1 antagonists and clinical characteristics of febrile seizures. *Int J Gen Med* 2012;5:277-281.
- Zuddas A, Zanni R, Usala T. Second generation antipsychotics (SGAs) for non-psychotic disorders in children and adolescents: a review of the randomized controlled studies. *Eur Neuropsychopharmacol* 2011;21:600-20.

Web Link References

21 CFR Part 201. Specific Requirements on Content and Format of Labeling for Human Prescription Drugs: Revision of "Pediatric Use" Subsection in the Labeling; Final Rule, Federal Register Volume 59, Number 238, December 13, 1994. <http://www.gpo.gov/fdsys/pkg/FR-1994-12-13/html/94-30238.htm>

Advisory Committee on Psychotropic Medications. The use of psychotropic medications for children and youth in the Texas foster care system. Texas Department of Family and Protective Services, September 1, 2004. Archived at: http://www.dfps.state.tx.us/ChildProtection/Medical_Services/guide-psychotropic.asp

Children and Adolescents' Psychoactive Medication Workgroup. Psychoactive medication for children and adolescents: Orientation for Parents, Guardians, and Others. Massachusetts Department of Mental Health, Boston, July 2007. <http://www.mass.gov/eohhs/docs/dmh/publications/psychoactive-booklet.pdf>

Child Exposure to Trauma: Comparative Effectiveness of Interventions Addressing Maltreatment (Review Number 89). Goldman FJ, Lloyd SW, et al., April 15, 2013. <http://effectivehealthcare.ahrq.gov/index.cfm/search-for-guides-reviews-and-reports/?pageaction=displayproduct&productid=1463>

Child Welfare Trauma Training Toolkit (2013). The National Child Traumatic Stress Network. <http://learn.nctsn.org/login/index.php>

Facts and Comparisons Drug Information. Clin-eguide [database online]. St. Louis, MO: Wolters Kluwer Health, Inc., 2012. <http://clineguide.ovid.com.ezproxy.lib.utexas.edu/>

FDA Drug Safety Communication: Revised recommendations for Celexa (citalopram hydrobromide) related to a potential risk of abnormal heart rhythms with high doses, August 24, 2011. <http://www.fda.gov/Drugs/DrugSafety/ucm297391.htm>

Health Care Issues for Children and Adolescents in Foster Care and Kinship Care. American Academy of Pediatrics Policy Statement, October 2015. <http://pediatrics.aappublications.org/content/136/4/e1131>

Making Healthy Choices: A Guide on Psychotropic Medication for Youth in Foster Care. Administration on Children, Youth and Families Children's Bureau, U.S. Department of Health and Human Services, 2012. <https://www.childwelfare.gov/pubs/makinghealthy-choices/>

Natural Medicines Comprehensive Database [database online]. Stockton, CA: Therapeutic Research Faculty, 2011. <http://naturaldatabase.therapeuticresearch.com>

Pediatric and Neonatal Lexi-Drugs. Lexi-Comp Online™ [database online]. Hudson, OH: Lexi-Comp, Inc., 2012. <http://online.lexi.com.ezproxy.lib.utexas.edu>

Recommendations about the Use of Psychotropic Medications for Children and Adolescents Involved in Child-Serving Systems. System of Care Resource from the American Academy of Child and Adolescent Psychiatry 2015. http://www.aacap.org/App_Themes/AACAP/docs/clinical_practice_center/systems_of_care/AACAP_Psychotropic_Medication_Recommendations_2015_FINAL.pdf

When to seek referral or consultation with a child or adolescent psychiatrist. American Academy of Child and Adolescent Psychiatry, 2003. http://www.aacap.org/AACAP/Member_Resources/Practice_Information/When_to_Seek_Referral_or_Consultation_with_a_CAP.aspx

Members of the Ad Hoc Working Group on Psychotropic Medication Parameters for Children and Youth in Foster Care

Chairs:

M. Lynn Crismon, PharmD: Psychopharmacologist, Dean, James T. Doluisio Regents Chair & Behrens Centennial Professor, College of Pharmacy, The University of Texas at Austin, Austin, TX.

James A. Rogers, MD: Child Psychiatrist, Medical Director, Texas Department of Family and Protective Services, Austin, TX.

Emilie Becker, MD: Child Psychiatrist, Medicaid/CHIP Mental Health Medical Director, Texas Health and Human Services Commission, Austin, TX.

Members:

Joseph Blader, PhD: Clinical Psychologist, Meadows Foundation & Semp Russ Professor of Child Psychiatry Research, Departments of Psychiatry and Pediatrics, The University of Texas Health Science Center at San Antonio, San Antonio, TX.

Angela Hughes Campbell, PharmD: Psychopharmacologist, Adjunct Assistant Professor, College of Pharmacy, The University of Texas at Austin, Nacogdoches, TX.

Mark Janes, MD: Child Psychiatrist, Medical Director, Bluebonnet Trails Community Services, Round Rock, TX.

Christopher J. Kratochvil, MD: Child Psychiatrist, Associate Vice Chancellor for Clinical Research, Chief Medical Officer UNeHealth, University of Nebraska Medical Center, Omaha, NE.

Molly Lopez, PhD: Clinical Psychologist, Research Associate Professor, School of Social Work, The University of Texas at Austin, Austin, TX.

James Lukefahr, MD: Pediatrician, Medical Director-Center for Miracles, Professor, Division of Child Abuse Pediatrics-The University of Texas Health Science Center, San Antonio, TX.

Octavio N. Martinez, Jr., MD: Psychiatrist, Executive Director-Hogg Foundation for Mental Health, The University of Texas at Austin, Austin, TX.

Nina Jo Muse, MD: Child Psychiatrist, Medical Director for Behavioral Health, Texas Department of State Health Services, Austin, TX.

Sylvia Muzquiz-Drummond, MD: Child Psychiatrist, Medical Director MHMRA of Harris County, Houston, TX.

Steven Pliszka, MD: Child Psychiatrist, Dielmann Distinguished Professor and Chair, Department of Psychiatry, The University of Texas Health Science Center, San Antonio, TX.

Dawnelle Schatte, MD: Child Psychiatrist, Associate Professor Department of Psychiatry and Behavioral Health Services, University of Texas Medical Branch, Galveston, TX.

Manuel Schydlower, MD: Pediatrician, Professor of Pediatrics, Associate Dean for Admissions, Texas Tech University Health Sciences Center, Paul L. Foster School of Medicine, El Paso, TX.

Observer/Monitors:

Amanda Garner, PharmD, Director of DUR/Formulary Management, Medicaid/CHIP Vendor Drug Program, Texas Health & Human Services Commission, Austin, TX.

Kay Lester, RPh, Pharmacy Benefit Oversight, Medicaid/CHIP Vendor Drug Program, Texas Health & Human Services Commission, Austin, TX

Committee Members Disclosures: Committee Disclosures: Since January 1, 2009, the authors below disclose the following financial relationships:

Dr. Blader has received funding as a consultant/researcher from Supernus Pharmaceuticals and research funding through his employer institution from Supernus.

Dr. Crismon has served on a one day advisory committee regarding biosimilars with funding through his employer institution from Amgen.

Dr. Kratochvil has received research funding through his employer institution from AstraZeneca, Abbott, Forest, Lilly, Neuren, Novartis, Pfizer, Otsuka, Seaside, Shire, and Somerset. He has received funding as an advisor through his employer institution from Abbott, AstraZeneca, Forest, Lilly, Pfizer, Seaside, and Shire Pharmaceuticals and, through his employer institution has received support for serving on the Data Safety Monitoring Boards for Neuren, Otsuka, Pfizer and Seaside Pharmaceuticals.

Dr. Lopez holds stock in Lilly, Merck, Proctor & Gamble, and Pfizer Pharmaceuticals.

Dr. Pliszka has received funding as a consultant for Ironshore and Shire Pharmaceuticals. Through his employer institution he has served as an expert witness for Eli Lilly and Janssen Pharmaceuticals. He has received research grants through his employer institution from Ironshore, Purdue and Shire.

The other members of the working group do not have any financial relationships to disclose.

Disclaimer

The authors of this document have worked to ensure that all information in the parameters is accurate at the time of publication and consistent with general psychiatric and medical standards and consistent with FDA labeling and information in the biomedical literature.

However, as medical research and practice continue to advance, therapeutic standards may change, and the clinician is encouraged to keep up with the current literature in psychiatry and clinical psychopharmacology. In addition, not all potential adverse drug reactions or complications are listed in the tables, and the clinician should consult the official FDA labeling and other authoritative reference sources for complete information.

These parameters are not a substitute for clinical judgement, and specific situations may require a specific therapeutic intervention not included in these parameters.

Acknowledgements

R. Joel Moore (Pharm.D. Candidate, The University of Texas at Austin) assisted with the literature search and updating of the medication tables.

Richard Steinberg (Accessibility Coordinator for Electronic and Information Resources, Texas Department of State Health Services) provided final editing and design.

Web Address for the ***March 2016 Psychotropic Medication Utilization Parameters for Children and Youth in Foster Care***

http://www.dfps.state.tx.us/Child_Protection/Medical_Services/guide-psychotropic.asp